

SEXUAL_HEALTH

The Importance of Regular STI Testing and How to Get Tested

Aruna Reddy*

* Correspondence: Aruna Reddy

Received: 27 January 2021

Published: 06 February 2021

Volume 8, Issue 1 (Jan–Mar 2021)

Article no.: e41

Open access · free to read

Introduction

Sexually Transmitted Infections (STIs) are a significant public health concern globally, affecting millions of individuals every year. Despite their prevalence, STIs often remain undiagnosed and untreated due to stigma, lack of awareness, and insufficient testing. Regular STI testing is a crucial component of sexual health, enabling early detection, treatment, and prevention of further transmission. This article aims to educate readers about the importance of regular STI testing, the process of getting tested, and the resources available for individuals seeking STI testing.

Understanding STIs

STIs are infections that are primarily spread through sexual contact, including vaginal, anal, and oral sex. There are more than 30 different bacteria, viruses, and parasites known to be transmitted through sexual contact. Common STIs include chlamydia, gonorrhea, syphilis, human immunodeficiency virus (HIV), human papillomavirus (HPV), and herpes simplex virus (HSV). STIs can have serious health consequences if left untreated, including infertility, chronic pain, and increased risk of acquiring other infections, such as HIV.

Types of STIs

STIs can be classified into three main categories:

- **Bacterial STIs:** These include chlamydia, gonorrhea, and syphilis. Bacterial STIs can usually be cured with antibiotics if detected early.
- **Viral STIs:** These include HIV, HPV, herpes, and hepatitis B. While viral STIs cannot be cured, they can be managed with medication to reduce symptoms and prevent transmission.
- **Parasitic STIs:** These include trichomoniasis and pubic lice (crabs). Parasitic STIs can be treated with prescription medication.

Why Regular STI Testing is Important

Regular STI testing is vital for maintaining sexual health and preventing the spread of infections. Here are some reasons why regular testing is crucial:

1. Early Detection and Treatment

Many STIs do not cause noticeable symptoms in their early stages, which means that individuals may be infected without knowing it. Regular testing allows for early detection, which is essential for prompt treatment. Early treatment can prevent complications and reduce the risk of transmitting the infection to others.

2. Reducing the Spread of Infections

Undiagnosed and untreated STIs can spread rapidly within communities, particularly among sexually active individuals. Regular testing helps identify and treat infections before they can be transmitted to others, thereby reducing the overall prevalence of STIs.

3. Protecting Reproductive Health

STIs can have serious consequences for reproductive health, particularly for women. Untreated STIs such as chlamydia and gonorrhea can lead to pelvic inflammatory disease (PID), which can cause infertility and chronic pelvic pain. Regular testing helps protect reproductive health by ensuring that infections are treated before they can cause long-term damage.

4. Reducing Stigma and Promoting Awareness

Regular STI testing can help reduce the stigma associated with STIs by normalizing the testing process. By making testing a routine part of sexual health care, individuals are more likely to feel comfortable discussing and addressing their sexual health concerns. Increased awareness and education about STIs can also encourage more people to get tested and seek treatment if needed.

Statistics on STI Prevalence

According to the World Health Organization (WHO), more than 1 million sexually transmitted infections are acquired every day worldwide. In the United States, the Centers for Disease Control and Prevention (CDC) estimates that nearly 20 million new STIs occur each year, with half of them affecting young people aged 15-24 years. Chlamydia and gonorrhea are among the most commonly reported STIs, with over 1.8 million cases of chlamydia reported in 2019 alone.

When to Get Tested

The frequency of STI testing depends on several factors, including age, sexual activity, and risk factors. Here are some general guidelines for STI testing:

1. Routine Testing

Sexually active individuals, particularly those with multiple partners, should consider getting tested at least once a year. Routine testing is also recommended for individuals in new sexual relationships, even if they do not have any symptoms.

2. After a New Sexual Partner

If you have recently had sex with a new partner, it is a good idea to get tested for STIs, especially if you did not use condoms or other barrier methods. Both partners should get tested before engaging in unprotected sex to ensure they are not unknowingly transmitting an infection.

3. Experiencing Symptoms

If you experience any symptoms of an STI, such as unusual discharge, burning during urination, sores or bumps on the genitals, or itching, you should get tested as soon as possible. Even if the symptoms go away on their own, it is still important to get tested to rule out an infection.

4. During Pregnancy

Pregnant women should be tested for STIs as part of their prenatal care. Some STIs can be passed from mother to baby during pregnancy or childbirth, so early detection and treatment are essential for protecting both the mother and the baby.

How to Get Tested for STIs

Getting tested for STIs is a straightforward process that can be done at various healthcare facilities. Here are the steps involved in getting tested:

1. Find a Testing Location

STI testing is available at many healthcare facilities, including:

- **Primary Care Clinics:** Many primary care providers offer STI testing as part of routine check-ups.
- **Sexual Health Clinics:** These clinics specialize in sexual health and often provide confidential STI testing.

- **Planned Parenthood:** Planned Parenthood clinics offer STI testing and treatment, along with other reproductive health services.
- **Community Health Centers:** Community health centers may offer free or low-cost STI testing for those without insurance.
- **Online Testing Services:** Some companies offer at-home STI testing kits that you can order online and mail back to a lab for results.

2. What to Expect During Testing

The testing process varies depending on the type of STI being tested for. Common testing methods include:

- **Urine Sample:** Used to test for chlamydia and gonorrhea.
- **Blood Test:** Used to test for HIV, syphilis, and hepatitis.
- **Swab Test:** A swab of the affected area (such as the genitals, throat, or rectum) may be taken to test for various STIs.
- **Physical Exam:** A healthcare provider may examine the genital area for signs of infection, such as sores or discharge.

Most STI tests are quick and painless, and results are usually available within a few days.

3. Understanding Your Results

After you receive your test results, it is important to discuss them with your healthcare provider. If you test positive for an STI, your provider will discuss treatment options with you. Many STIs are curable with medication, and those that are not curable can be managed with treatment to reduce symptoms and prevent transmission. If you test negative, your provider may recommend future testing based on your risk factors and sexual activity.

4. Partner Notification

If you test positive for an STI, it is important to notify your sexual partners so they can also get tested and treated if necessary. Many healthcare providers offer confidential partner notification services, where they can anonymously inform your partners without revealing your identity.

Preventing STIs

While regular testing is important, prevention is the best way to protect yourself and your partners from STIs. Here are some strategies for reducing your risk of contracting an STI:

1. Use Condoms Consistently

Condoms are highly effective at preventing the transmission of most STIs, including HIV, chlamydia, and gonorrhea. Use condoms consistently and correctly every time you have vaginal, anal, or oral sex.

2. Get Vaccinated

Vaccines are available to protect against certain STIs, including HPV and hepatitis B. Talk to your healthcare provider about getting vaccinated if you are eligible.

3. Limit Your Number of Sexual Partners

The more sexual partners you have, the greater your risk of being exposed to an STI. Limiting your number of partners can reduce your risk of infection.

4. Get Tested Regularly

Regular STI testing is essential for early detection and treatment. Make testing a routine part of your sexual health care.

5. Communicate with Your Partners

Open and honest communication with your sexual partners is key to maintaining sexual health. Discuss STI testing, condom use, and sexual history with your partners before engaging in sexual activity.

Conclusion

Regular STI testing is a critical component of sexual health that helps prevent the spread of infections, protects reproductive health, and reduces stigma. By understanding the importance of testing and knowing how to get tested, individuals can take proactive steps to protect themselves and their partners. Remember, sexual health is an important part of overall well-being, and regular testing is a simple and effective way to ensure that you stay healthy.

Competing interests

The authors declare that they have no competing interests.

How to cite this article (Vancouver)

Reddy A. The Importance of Regular STI Testing and How to Get Tested. CHEARS Chronical. 2021;8(1):e41. Available from: <https://chears.in/journal/volume-8/issue-1/the-importance-of-regular-sti-testing-and-how-to-get-tested-41>

© 2021 The authors. Published by CHEARS in CHEARS Chronical.

<https://chears.in/journal/volume-8/issue-1/the-importance-of-regular-sti-testing-and-how-to-get-tested-41>